



## Intraday effects of ambient PM<sub>1</sub> on emergency department visits in Guangzhou, China: A case-crossover study

Linjiong Liu<sup>a</sup>, Fujian Song<sup>b</sup>, Jiaying Fang<sup>c</sup>, Jing Wei<sup>d,e</sup>, Hung Chak Ho<sup>f</sup>, Yimeng Song<sup>f</sup>, Yuanyuan Zhang<sup>a</sup>, Lu Wang<sup>g</sup>, Zhiming Yang<sup>h</sup>, Chengyang Hu<sup>i,j</sup>, Yunquan Zhang<sup>a,k,\*</sup>

<sup>a</sup> Department of Epidemiology and Biostatistics, School of Public Health, Medical College, Wuhan University of Science and Technology, Wuhan 430065, China

<sup>b</sup> Public Health and Health Services Research, Norwich Medical School, University of East Anglia, Norwich, NR4 7TJ, UK

<sup>c</sup> Huadu District People's Hospital, Southern Medical University, Guangzhou 510800, China

<sup>d</sup> State Key Laboratory of Remote Sensing Science, College of Global Change and Earth System Science, Beijing Normal University, Beijing, China

<sup>e</sup> Department of Atmospheric and Oceanic Science, Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD, USA

<sup>f</sup> Department of Urban Planning and Design, The University of Hong Kong, Hong Kong, China

<sup>g</sup> Department of Nursing, Medical College, Wuhan University of Science and Technology, Wuhan 430065, China

<sup>h</sup> Donlinks School of Economics and Management, University of Science and Technology Beijing, Beijing 100083, China

<sup>i</sup> Department of Humanistic Medicine, School of Humanistic Medicine, Anhui Medical University, Hefei 230032, China


<sup>j</sup> Department of Epidemiology and Biostatistics, School of Public Health, Anhui Medical University, Hefei 230032, China

<sup>k</sup> Hubei Province Key Laboratory of Occupational Hazard Identification and Control, Wuhan University of Science and Technology, Wuhan 430065, China

### HIGHLIGHTS

- First study to assess detrimental effects of PM<sub>1</sub> at a sub-daily timescale
- Exposure to ambient PMs will elevate risks of ED visits in a few hours.
- PM<sub>1</sub> can induce slightly stronger effects compared with PM<sub>2.5</sub> and PM<sub>10</sub>.
- Significantly more potent PM-associated effects during the cold months

### GRAPHICAL ABSTRACT



### ARTICLE INFO

#### Article history:

Received 26 June 2020

Received in revised form 6 September 2020

Accepted 9 September 2020

Available online 15 September 2020

Editor: Pavlos Kassomenos

#### Keywords:

PM<sub>1</sub>

Hourly effects

Emergency department visits

Case-crossover design

China

### ABSTRACT

**Background:** Short-term exposure to PM<sub>2.5</sub> has been widely associated with human morbidity and mortality. However, most up-to-date research was conducted at a daily timescale, neglecting the intra-day variations in both exposure and outcome. As an important fraction in PM<sub>2.5</sub>, PM<sub>1</sub> has not been investigated about the very acute effects within a few hours.

**Methods:** Hourly data for size-specific PMs (i.e., PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub>), all-cause emergency department (ED) visits and meteorological factors were collected from Guangzhou, China, 2015–2016. A time-stratified case-crossover design with conditional logistic regression analysis was performed to evaluate the hourly association between size-specific PMs and ED visits, adjusting for hourly mean temperature and relative humidity. Subgroup analyses stratified by age, sex and season were conducted to identify potential effect modifiers.

**Results:** A total of 292,743 cases of ED visits were included. The effects of size-specific PMs exhibited highly similar lag patterns, wherein estimated odds ratio (OR) experienced a slight rise from lag 0–3 to 4–6 h and subsequently attenuated to null along with the extension of lag periods. In comparison with PM<sub>2.5</sub> and PM<sub>10</sub>, PM<sub>1</sub> induced slightly larger effects on ED visits. At lag 0–3 h, for instance, ED visits increased by 1.49% (95% confidence

\* Corresponding author at: Department of Epidemiology and Biostatistics, School of Public Health, Medical College, Wuhan University of Science and Technology, Wuhan 430065, China.  
E-mail address: [YunquanZhang@wust.edu.cn](mailto:YunquanZhang@wust.edu.cn) (Y. Zhang).

interval: 1.18–1.79%), 1.39% (1.12–1.66%) and 1.18% (0.97–1.40%) associated with a 10- $\mu\text{g}/\text{m}^3$  rise, respectively, in  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$ . We have detected a significant effect modification by season, with larger  $\text{PM}_{10}$ -associated OR during the cold months (1.017, 1.013 to 1.021) compared with the warm months (1.010, 1.005 to 1.015).

**Conclusions:** Our study provided brand-new evidence regarding the adverse impact of  $\text{PM}_{10}$  exposure on human health within several hours.  $\text{PM}_{10}$ -associated effects were significantly more potent during the cold months. These findings may aid health policy-makers in establishing hourly air quality standards and optimizing the allocation of emergency medical resources.

© 2020 Elsevier B.V. All rights reserved.

## 1. Introduction

Ambient particulate matter has been linked to adverse health outcomes, including morbidity and mortality (Liu et al., 2019; Tian et al., 2019b). Reported by *State of Global Air 2019*, ambient particulate matter with an aerodynamic diameter less than 2.5  $\mu\text{m}$  ( $\text{PM}_{2.5}$ ) contributed to nearly three million deaths in 2017, and more than half of this disease burden fell on densely populated countries, such as China and India (Institute, 2019). China has been experiencing severe particulate pollution, particularly in metropolises such as Beijing and Guangzhou (Guo et al., 2016; Guo et al., 2018; Zhou et al., 2019). In China,  $\text{PM}_{2.5}$  pollution ranked fourth in 2017 as a risk factor for both the magnitude of deaths and disability adjusted life years (DALYs) (Zhou et al., 2019). A large-scale modeling study in China estimated that 0.14 and 0.06 years of gain in life expectancy could be achieved by reducing concentrations of ambient  $\text{PM}_{2.5}$  to 25 and 75  $\mu\text{g}/\text{m}^3$ , respectively (Qi et al., 2020).

Numerous epidemiological studies found that long- and short-term exposures to ambient particulate matter with an aerodynamic diameter less than 10  $\mu\text{m}$  ( $\text{PM}_{10}$ ) or  $\text{PM}_{2.5}$  are closely associated with increased morbidity and mortality, including but not limited to cardiovascular and respiratory diseases (Bhaskaran et al., 2011; Huang et al., 2019; Hvidtfeldt et al., 2019; Yin et al., 2020). For the short-term effects of  $\text{PM}_{10}$ , emergency department (ED) visits have been frequently used as a surrogate of acute disease events (Yorifuji et al., 2014a; Zhang et al., 2019b). Other proxies include hospital admissions (Liu et al., 2018), emergency ambulance calls (Ai et al., 2019) and so forth. However, most existing studies were conducted at a daily timescale, without considering the intra-day variations in both exposure and outcome (Lin et al., 2017b). In recent years, hourly associations between exposure to ambient fine and inhalable  $\text{PM}_{10}$  and human morbidity have been received increasing attention, being focused on outcomes of hospital admission for cardiovascular and respiratory diseases (Bhaskaran et al., 2012; Chen et al., 2020a; Kim et al., 2015), as well as ambulance calls and ED visits (Ai et al., 2019; Chen et al., 2019a; Phung et al., 2020). Nevertheless, particulate matter with an aerodynamic diameter less than 1  $\mu\text{m}$  ( $\text{PM}_1$ ) has not been investigated in the aforementioned studies. Additionally, few studies investigated the associations between  $\text{PM}_1$  and health outcomes at other longer timescales (e.g., daily), probably due to the lack of routine surveillance and very sparse data from ground measurements of  $\text{PM}_1$  (Chen et al., 2018; Zhang et al., 2019b).

Compared with larger fractions,  $\text{PM}_1$  has higher surface-to-volume ratios, greater vascular penetration, and contains more toxins (Chen et al., 2017; Chen et al., 2019b; Yang et al., 2019). It is noteworthy that, consistently high  $\text{PM}_1/\text{PM}_{2.5}$  ratios (i.e., range from 0.75 to 0.88) were reported across China (Chen et al., 2018; Guo et al., 2009; Zang et al., 2019), suggesting that  $\text{PM}_1$  is a crucial driver of  $\text{PM}_{2.5}$  pollution and accounts for a large proportion of  $\text{PM}_{2.5}$  (Wang et al., 2019; Zang et al., 2018). Another study in China also supported the notion that smaller size fractions of  $\text{PM}_{10}$  have more detrimental mortality impacts (Hu et al., 2018). Nevertheless, the influential mechanisms have yet to be well understood, especially for the associations of  $\text{PM}_1$  with human health, which merits further analyses.

To fill the research gap, this study aimed to investigate the hourly associations between emergency department (ED) visits and exposures to ambient size-specific  $\text{PM}_{10}$ , particularly  $\text{PM}_1$ , by adopting a time-

stratified case-crossover design. Subgroup analyses stratified by age group and sex were conducted to identify potentially vulnerable sub-populations. We also assessed seasonal patterns of  $\text{PM}_{10}$ -related effects by dividing the whole study period into warm (April to September) and cold (October to March of next year) months.

## 2. Materials and methods

### 2.1. Study setting and population

Guangzhou, the capital of Guangdong province, China, has a typical subtropical humid monsoon climate. Because of the rapid development of economy and the increment of energy consumption during the past several decades, Guangzhou has been affected by severe air pollution. Huadu District People's Hospital is a large-scale tertiary general hospital, situated in an urban area of Huadu District in Guangzhou. In 2018, over one million permanent residents lived in this district (<http://tjj.gz.gov.cn/>). Our study included patients who presented to the emergency department (ED) of this hospital during 2015–2016.

### 2.2. Data collection

Individual records of all-cause emergency department visit through January 1, 2015 to December 31, 2016 were collected from Huadu District People's Hospital of Guangzhou, China. The primary characteristics of each case were extracted, including the specific time of admission, sex and age. We aggregated hourly counts of all-cause ED visits as the health outcome at each calendar day during our study period.

Ground-based measurements of hourly mean concentrations of ambient size-specific  $\text{PM}_{10}$  (i.e.,  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$ , and  $\text{PM}_{10}$ ) during 2015–2016 were obtained from Atmosphere Watch Network (CAWNET), run by the China Meteorological Administration (Chen et al., 2018). Due to the lack of routine surveillance and the sparse monitoring stations of  $\text{PM}_1$  in China, our  $\text{PM}_1$  data were collected from two fixed stations which can monitor  $\text{PM}_1$  in Guangzhou (Wei et al., 2019a). In order to facilitate comparability of risk assessments between size-specific  $\text{PM}_{10}$ , we also used the same stations' data of  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  for exposure assessments. Owing to data unavailability of residential addresses for ED patients, we averaged these  $\text{PM}_{10}$  measurements from two fixed stations as the substitute of population-level particulate exposure in our analysis. At the same period, we also collected hourly average concentrations of gaseous pollutants (i.e.,  $\text{SO}_2$ ,  $\text{NO}_2$ ,  $\text{O}_3$  and CO) from the China National Urban Air Quality Real-time Publishing Platform (<http://106.37.208.233:20035/>), and hourly meteorological data (i.e., mean temperature and relative humidity) from the global hourly datasets of United States' National Centers for Environmental Information (NCEI, <https://www.ncei.noaa.gov/>).

In addition, we collected hourly precipitation data for Guangzhou, 2015–2016, from the ERA-5 reanalysis dataset on the Copernicus climate data store (ECMWF, ERA5 hourly data on single levels from 1979 to present, <https://cds.climate.copernicus.eu>). Then we transformed hourly precipitation data into a binary variable of rains (whether it rains or not in a specific hour).

Final dataset for analysis was comprised of these hourly ambient data and ED visits matched through the corresponding time windows

(i.e., the same hour or hours). Due to monitor network system issues, 748 (4.3%) and 733 (4.2%) hours' data were not registered for size-specific PMs and gaseous pollutants. In our analysis, we thus excluded cases of ED visits during these exposure windows.

### 2.3. Statistical analyses

We adopted a time-stratified case-crossover (TSCC) design to separately investigate the hourly association between size-specific PMs (i.e., PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub>) exposure and ED visits. Case-crossover design can be deemed to be an extension of the traditional case of matched case-control design, which incorporates the advantages of both case-control and cross-sectional study (Maclure, 1991). Time-stratified case-crossover (TSCC) design was proposed by Lumley and Levy in 2000 based on general case-crossover design, which used the day or days within the same time tier as the control group to control exposures (Lumley and Levy, 2000). TSCC design was widely employed to investigate the short-term effects of environmental factors (e.g., air pollution and extreme weather conditions) on adverse health outcomes, including morbidity and mortality (Bhaskaran et al., 2011; Di et al., 2017; Yorifuji et al., 2014a). In this study design, all cases act as their own controls meanwhile we used calendar month and year as fixed time strata, which can effectively control long-term trends and seasonality and effects of time-invariant individual-level confounders such as age, sex, behavior and metabolic factors (Lumley and Levy, 2000; Wei et al., 2019b). Moreover, to avoid impacts from days of the week and intra-day variation, we selected control periods from the same hour of the same day of the week in the calendar month of ED visits (Chen et al., 2020a; Phung et al., 2020). For each case, we assigned three or four matched controls that have not occurred of ED visits at the timescale.

A conditional logistic regression (CLR) was used to fit TSCC model and separately assess the associations between transient exposures to size-specific particulate matters and ED visits (Di et al., 2017; Wei et al., 2019b). In our analytical models, we used natural cubic spline (NS) function with three degrees of freedom (df) to fit the effects of hourly current temperature (Temp<sub>0</sub>) and relative humidity (Rh<sub>0</sub>). Then we introduced them as covariate terms to eliminate nonlinear confounding effects (Jiao et al., 2020; Zhang et al., 2019b). In this study, the conditional logistic regression model was implemented through Cox proportional hazard regression method (Le and Lindgren, 1988) and presented as below:  $\ln[h(t,X)] = \ln[h_0(t)] + \beta(PMs) + NS(Temp_0, df = 3) + NS(Rh_0, df = 3)$ , where  $t$  refers to the specific hour of interest;  $X$  refers to the record of ED visit;  $h_0(t)$  and  $h(t,X)$  represents the baseline and estimated hazard function at hour  $t$ , respectively;  $\beta$  refers to estimated coefficient for PMs and PMs refers to hourly mean concentrations of PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub>;  $NS(Temp_0)$  and  $NS(Rh_0)$  are NS functions for Temp<sub>0</sub> and Rh<sub>0</sub>, respectively.

Meanwhile, we examined the linearity hypothesis between size-specific PMs and ED visits by smoothing exposure-response curves using NS terms with three df for PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub>. To better capture the very short-term effects on health from size-specific PMs, we divided exposure time windows into a set of lag periods (e.g., 0–3 h, 4–6 h and 7–12 h) up to 72 h before the event of ED visits (Kim et al., 2015). For instance, lag 0–3 h means the moving average concentration of 4 h before the event of interest. Moreover, these lag periods were selected as a compromise between model parsimony and flexibility. In this study, the risk of ED visits was estimated and reported as odds ratio (OR) and corresponding 95% confidence interval (CI) associated with a 10- $\mu\text{g}/\text{m}^3$  increase in PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> concentrations at various lag periods, respectively.

Besides, we performed a string of subgroup analyses stratified by sex (i.e., male and female) and age (i.e., 0–14, 15–34, 35–64 and over 65 years) to identify potential susceptible subpopulations. These age strata comply with the age distribution of the study population and some previous literature (Jiao et al., 2020; Zhang et al., 2019b). Also, seasonal analyses of PM-associated effects were conducted by dividing the

whole study period into warm (April to September) and cold (October to March of next year) months. In these subgroup analyses, a two-sample Z test (Zhang et al., 2020) and meta-regression methods (Guo, 2017; Tian et al., 2019b) were applied to assess whether age, sex, and season were sources of potential modification effects.

### 2.4. Sensitivity analyses

To verify the robustness of our findings, we changed the parameter specifications in our modeling strategies by (1) performing a two-pollutant analysis through separately including gaseous pollutants (i.e., SO<sub>2</sub>, NO<sub>2</sub>, O<sub>3</sub> and CO), (2) treating whether it rains or not in a specific hour as a binary covariate and including it in our main model, and (3) varying dfs from 3 to 6 for NCS function terms and exposure periods of meteorological factor (i.e., from lag 0 to lags 0–3 and 0–6 h). Moreover, we used the aforementioned meta-regression method to test the statistical significance of difference before and after adjustment.

All data analyses were run on R software (version 3.6.2). We used the "survival" package for conditional logistic regression modeling and the "mvmeta" package for meta-regression analysis. Two-sided tests with a  $p$ -value of less than 0.05 were considered to be statistically significant.

## 3. Results

Table 1 describes the basic characteristics of all-cause ED patients. A total of 292,743 cases were involved, with hourly mean visits of 16.7 (standard deviation [SD], 12.1). Approximately 53.3% of patients were male and over two-thirds were young people aged less than 35 years old. Relatively more ED visits occurred in the warm months and accounted for 53.4% of total cases. Characteristics in seasonal patterns of ED visits were summarized in Table S1, stratified by sex and age group.

Table 2 outlines the distribution characteristics of ambient air pollution and meteorological factors during 2015–2016 in Guangzhou. Hourly mean (SD) concentrations were 26.7 (16.4)  $\mu\text{g}/\text{m}^3$  for PM<sub>1</sub>, 31.0 (18.3)  $\mu\text{g}/\text{m}^3$  for PM<sub>2.5</sub> and 40.2 (23.4)  $\mu\text{g}/\text{m}^3$  for PM<sub>10</sub>, respectively. Annual average temperature and relative humidity was 23.4 °C (Range: 2.5–37.5 °C) and 74.9% (Range: 16.5–100%), respectively. Ambient levels of all pollutants were higher during the cold months, except for ozone (Table S2). Fig. 1 shows high correlations between size-fractional PMs, with Spearman correlation coefficients ranging from 0.93 to 0.98. Conversely, PMs were low-to-moderate correlated with gaseous pollutants and climate variables.

Fig. 2 manifests dose-response associations of all-cause ED visits with PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub>, respectively. Size-fractional PMs' curves share approximate linearity and similar pattern with a steeper slope

**Table 1**  
Basic characteristics of emergency department visits in Guangzhou, China, 2015–2016.

Group	Total		Hourly	
	Count (n)	Percentage (%)	Mean	SD
All	292,793	100.0	16.7	12.1
Sex				
Male	161,953	55.3	9.2	6.8
Female	130,840	44.7	7.5	6.0
Age, years				
<15	97,508	33.3	5.6	5.5
15–34	90,847	31.0	5.1	4.4
35–64	79,796	27.3	4.5	3.4
≥65	24,642	8.4	1.4	1.5
Season				
Warm	156,247	53.4	17.8	12.5
Cold	136,546	46.6	15.9	11.5

Abbreviations: SD, standard deviation; Warm season, April to September; Cold season, October to March of the next year.

**Table 2**  
Hourly distributions in ambient air pollution and meteorological factors during 2015–2016 in Guangzhou.

Variable	Mean	SD	Min	Percentiles			Max
				25th	50th	75th	
<b>Particulate pollutants</b>							
PM <sub>1</sub> , µg/m <sup>3</sup>	26.7	16.4	0.6	14.1	23.5	35.4	121.5
PM <sub>2.5</sub> , µg/m <sup>3</sup>	31.0	18.3	1.0	17.1	27.3	40.7	164.1
PM <sub>10</sub> , µg/m <sup>3</sup>	40.2	23.4	1.1	22.7	35.3	53.1	223.7
<b>Gaseous pollutants</b>							
NO <sub>2</sub> , µg/m <sup>3</sup>	43.9	23.1	4.0	28.0	39.0	53.0	225.0
SO <sub>2</sub> , µg/m <sup>3</sup>	11.7	6.7	2.0	7.0	10.0	15.0	106.0
O <sub>3</sub> , µg/m <sup>3</sup>	43.3	44.8	1.0	12.0	27.0	58.0	334.0
CO, mg/m <sup>3</sup>	1.0	0.3	0.4	0.8	0.9	1.1	3.0
<b>Weather conditions</b>							
Temperature, °C	23.4	6.7	2.5	18.5	25.0	28.0	37.5
Relative humidity, %	74.9	16.5	16.5	64.0	77.1	88.5	100.0

Abbreviations: SD, standard deviation; PM<sub>1</sub>, particulate matter with aerodynamic diameter ≤ 1 µm; PM<sub>2.5</sub>, particulate matter with aerodynamic diameter ≤ 2.5 µm; PM<sub>10</sub>, particulate matter with aerodynamic diameter ≤ 10 µm; NO<sub>2</sub>, nitrogen dioxide; SO<sub>2</sub>, sulfur dioxide; O<sub>3</sub>, ozone; CO, carbon monoxide.

in the low concentrations. As illustrated in Fig. 3, transient exposures to PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> were significantly associated with increased risks of ED visits in the ensuing 48 h. Also, the effects of size-specific PMs exhibited highly similar lag patterns, wherein estimated ORs experienced a slight rise from lag 0–3 to 4–6 h and subsequently attenuated to null along with the extension of lag periods. In comparison with PM<sub>2.5</sub> and PM<sub>10</sub>, PM<sub>1</sub> induced slightly larger effects on ED visits. With a 10-µg/m<sup>3</sup> rise in PM exposure, for instance, risks at lag 0–3 h increased by 1.49% (95% CI: 1.18–1.79%) for PM<sub>1</sub>, 1.39% (1.12–1.66%) for PM<sub>2.5</sub>, and 1.18% (0.97–1.40%) for PM<sub>10</sub>, respectively. Estimates for PM-associated ORs across various lag periods were detailed in Table S3.

Fig. 4 shows OR estimates of subgroups stratified by sex and age. We identified remarkable associations between PM and ED visits in all


subgroups, while no significant differences existed between genders as well as age groups. Specifically, ORs associated with a 10-µg/m<sup>3</sup> rise in PM<sub>1</sub> were comparable between males (1.016, 1.012 to 1.020) and females (1.014, 1.009 to 1.018), but were generally larger among children (1.018, 1.012 to 1.023) than older groups. Similar findings were also observed in associations of ED visits with PM<sub>2.5</sub> and PM<sub>10</sub>.

Table 3 gives season-specific OR estimates at two specific lag periods of lag 0–3 and 0–6 h, associated with per 10-µg/m<sup>3</sup> increase in PM exposure. In total population, significant effects from exposure to PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> were found in both warm and cold months. However, warm-cold differences in PM effects were solely evident in PM<sub>1</sub> at lag 0–3 h (*p*-value = 0.033), with a stronger association in the cold months (OR = 1.017, 95% CI: 1.013 to 1.021) than in the warm months (1.010, 1.005 to 1.015). Furthermore, in age groups except for children, the effects of hourly PMs on ED visits mostly exist in the cold months (Table S4). Compared with the all-age population, seasonal differences were more conspicuous in the age group of 15–34 years, with insignificance (*p* = 0.061) only occurring in PM<sub>10</sub> at lag 0–6 h. More details for seasonal analyses stratified by sex and age can be found in Table S4.


Sensitivity analysis demonstrates the robustness of our main findings. The OR estimates changed little (e.g., ranging from 1.015 to 1.014 for PM<sub>1</sub>) when varying dfs and lag periods for temperature and humidity (Table S5). In our two-pollutant modeling analyses, the estimated associations did not change dramatically except for adjusting NO<sub>2</sub>. For instance, PM<sub>1</sub>-associated ORs significantly decreased from 1.015 (1.012 to 1.018) to 1.006 (1.002 to 1.010) after including NO<sub>2</sub> in the two-pollutant model.

#### 4. Discussion

To the best of our knowledge, this is the first study concurrently assessing the associations between exposure to PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> and all-cause ED visits at a sub-daily timescale. In this study, we



**Fig. 1.** Spearman correlation matrix between ambient air pollutants and meteorological factors in Guangzhou, China, 2015–2016. Abbreviations: PM<sub>1</sub>, particulate matter with aerodynamic diameter ≤ 1 µm; PM<sub>2.5</sub>, particulate matter with aerodynamic diameter ≤ 2.5 µm; PM<sub>10</sub>, particulate matter with aerodynamic diameter ≤ 10 µm; SO<sub>2</sub>, sulfur dioxide; NO<sub>2</sub>, nitrogen dioxide; CO, carbon monoxide; O<sub>3</sub>, ozone; Temp, temperature; Rh, relative humidity.




**Fig. 2.** Exposure-response curves for PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> associated with all-cause ED visits at lag 0–3 h, respectively. Abbreviations: ED, emergency department; PM<sub>1</sub>, particulate matter with aerodynamic diameter ≤ 1 μm; PM<sub>2.5</sub>, particulate matter with aerodynamic diameter ≤ 2.5 μm; PM<sub>10</sub>, particulate matter with aerodynamic diameter ≤ 10 μm.


employed a time-stratified case-crossover design to evaluate acute effects from specific-PMs on human health. We found that transient exposures to ambient PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> significantly increased risks of ED visits in a few hours. Seasonal modification effects were detected in subgroup analyses, with PM<sub>1</sub>-associated OR estimates substantially larger during the cold months.

Over the past years, some researchers reported hourly associations between exposures to ambient fine and inhalable PMs and human morbidity, being focused on outcomes of hospital admission for cardiovascular and respiratory diseases (Bhaskaran et al., 2012; Chen et al., 2020a; Kim et al., 2015), as well as ambulance calls and ED visits (Ai et al., 2019; Chen et al., 2019a; Phung et al., 2020). However, PM<sub>1</sub>, a smaller particulate matter, has not been taken into account in the studies mentioned above. In our study, through using hourly mean concentrations of size-specific PMs as independent variables in analytic models, we incorporated the information of variation within a day or even a few hours.

Our results showed that increased all-cause ED visits were significantly associated with exposures to PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> within a few hours. Moreover, a mild and transient ascending trend in PM-associated ORs was observed from lag 0–3 to 4–6 h, followed by a descending trend at longer lags, suggesting that the first few hours may be the critical exposure window. These significant associations of interest remained up to 48 h after exposures. Our findings were echoed with a case-crossover study of eleven cities in Japan (Phung et al., 2020) and another study in Shenzhen city, China (Chen et al., 2019a). The former study (Phung et al., 2020) revealed that transient exposure to ambient PM<sub>2.5</sub> in a few hours might trigger elevated all-cause emergency ambulance dispatches, and their results manifested that these effects remained significant until to 24 h later. Furthermore, hourly associations between PMs (i.e., PM<sub>2.5</sub> and PM<sub>10</sub>) and ED visits persisted for about 10 h before the event of interest in the latter study (Chen et al., 2019a). Compared with the two studies above, our findings manifest longer sustained effects from specific-PMs on acute disease events,



**Fig. 3.** Odds ratios (with 95% CIs) for all-cause ED visits in Guangzhou at various exposure hours, associated with per 10 μg/m<sup>3</sup> increase in PM<sub>1</sub> (a), PM<sub>2.5</sub> (b), and PM<sub>10</sub> (c). Notes: \**p* < 0.05; \*\**p* < 0.001. Abbreviations: CI, confidence interval; ED, emergency department; PM<sub>1</sub>, particulate matter with aerodynamic diameter ≤ 1 μm; PM<sub>2.5</sub>, particulate matter with aerodynamic diameter ≤ 2.5 μm; PM<sub>10</sub>, particulate matter with aerodynamic diameter ≤ 10 μm.



**Fig. 4.** Odds ratios (with 95% CIs) for ED visits in Guangzhou among subgroups stratified by sex and age, associated with per 10  $\mu\text{g}/\text{m}^3$  increase in exposure to PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub> at lag 0–3 h. Notes: \* $p < 0.05$ ; \*\* $p < 0.001$ ; <sup>a</sup> p-value for difference between subgroups. Abbreviations: CI, confidence interval; ED, emergency department; PM<sub>1</sub>, particulate matter with aerodynamic diameter  $\leq 1 \mu\text{m}$ ; PM<sub>2.5</sub>, particulate matter with aerodynamic diameter  $\leq 2.5 \mu\text{m}$ ; PM<sub>10</sub>, particulate matter with aerodynamic diameter  $\leq 10 \mu\text{m}$ .

which might be explained by different demographic characteristics and pollutant levels. Nevertheless, the overall conclusion is still consistent, suggesting that daily air quality guidelines based on mean concentrations of air pollutants may be inadequate to protect human health (Lin et al., 2017a; Yorifuji et al., 2014b). Relevant studies are warranted in the future as references for the appropriate temporal scale of air quality standards.

The exposure-response relationships have a monotonic increasing trend, while we observed the steeper slope in low PM concentrations. This finding suggested that people may be more sensitive to adverse effects from ambient PMs at low levels, which is echoed with previous studies. A large time-series study in 184 major Chinese cities (Tian et al., 2019b) found that the dose-response curve increased sharply at low PM<sub>2.5</sub> concentrations without a discernible threshold. Another study (Liu et al., 2019) also reported that both curves were steeper at levels lower than 20  $\mu\text{g}/\text{m}^3$  for PM<sub>2.5</sub> and 40  $\mu\text{g}/\text{m}^3$  for PM<sub>10</sub>. More

investigations on hourly associations between ambient PM and human health are needed, which may provide a valuable reference for establishing hourly air quality limits.

Compared with PM<sub>2.5</sub> and PM<sub>10</sub>, PM<sub>1</sub> has been considered with more detrimental impacts on human health, likely owing to the higher surface-to-volume ratios, greater vascular penetration, and more toxins (Chen et al., 2017; Franck et al., 2011; Wu et al., 2020; Yin et al., 2020). A recent nationwide study (Yin et al., 2020) found that PM<sub>1</sub>-related excess CVD risk (0.29%, 95%CI: 0.12 to 0.47%) was significantly higher than PM<sub>2.5</sub> (0.24%) and PM<sub>10</sub> (0.21%). A multi-city study (Chen et al., 2017) also supported this notion and compared the effects of specific-PMs (i.e., PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>1–2.5</sub>) on ED visits. Their findings suggested that most of ED visits attributed to PM<sub>2.5</sub> are accounted for PM<sub>1</sub>, in that no associations between PM<sub>1–2.5</sub> and ED visits. In the present study, our comparative analyses also showed that smaller PM fraction exhibited more adverse effects on risks of ED visits. Specifically, risks

**Table 3**  
Season-specific ORs [95% CIs] of ED visits at lag 0–3 and 0–6 h, associated with per 10- $\mu\text{g}/\text{m}^3$  increase in PM exposures.

Group	Season	0–3 h		0–6 h		
		OR [95% CI]	P for interaction	OR [95% CI]	P for interaction	
All	PM <sub>1</sub>	Cold	1.017 [1.013 to 1.021]**	0.033	1.017 [1.013 to 1.021]**	0.053
		Warm	1.010 [1.005 to 1.015]**		1.011 [1.006 to 1.016]**	
	PM <sub>2.5</sub>	Cold	1.015 [1.011 to 1.018]**	0.103	1.015 [1.012 to 1.019]**	0.183
		Warm	1.010 [1.006 to 1.014]**		1.012 [1.007 to 1.016]**	
	PM <sub>10</sub>	Cold	1.013 [1.010 to 1.015]**	0.109	1.013 [1.010 to 1.016]**	0.265
		Warm	1.009 [1.006 to 1.012]**		1.011 [1.007 to 1.014]**	
Age 15–34	PM <sub>1</sub>	Cold	1.021 [1.014 to 1.028]**	0.007	1.021 [1.014 to 1.028]**	0.016
		Warm	1.006 [0.998 to 1.015]		1.007 [0.998 to 1.016]	
	PM <sub>2.5</sub>	Cold	1.019 [1.013 to 1.025]**	0.019	1.019 [1.012 to 1.025]**	0.036
		Warm	1.007 [0.999 to 1.015]		1.008 [1.000 to 1.016]*	
	PM <sub>10</sub>	Cold	1.015 [1.010 to 1.020]**	0.039	1.015 [1.010 to 1.020]**	0.061
		Warm	1.007 [1.001 to 1.013]*		1.008 [1.002 to 1.014]*	

Abbreviations: OR, odds ratio; CI, confidence interval; ED, emergency department; PM<sub>1</sub>, particulate matter with aerodynamic diameter  $\leq 1 \mu\text{m}$ ; PM<sub>2.5</sub>, particulate matter with aerodynamic diameter  $\leq 2.5 \mu\text{m}$ ; PM<sub>10</sub>, particulate matter with aerodynamic diameter  $\leq 10 \mu\text{m}$ ; Warm season, April to September; Cold season, October to March of the next year.

\*  $p < 0.05$ .

\*\*  $p < 0.001$ .

for ED visits at lag 0–3 h increased by 1.49% (95% CI: 1.18–1.79%), 1.39% (1.12–1.66%) and 1.18% (0.97–1.40%) associated with a 10- $\mu\text{g}/\text{m}^3$  increment in PM<sub>1</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> concentrations, respectively. Previously proposed biological mechanisms for the associations between particulate matter and acute disease events involved PM-augmented systematic inflammation and oxidative stress (Chen et al., 2020a; Lin et al., 2016; Yang et al., 2018). More mechanism researches are recommended to elucidate the exact biological pathway about the associations between ambient PM and acute disease events, particularly in smaller matters (e.g., PM<sub>1</sub> and ultrafine particles).

Prior evidence about PM effects modified by age and sex were mixed. In the present study, we observed comparable PM-associated effects between genders, in line with a nationwide time-series study (Tian et al., 2019a) on adult hospital admissions in 200 Chinese cities. However, another multi-city study (Chen et al., 2017) on daily ED visits reported higher relative risks of PM<sub>1</sub> and PM<sub>2.5</sub> on women. Kan and colleagues (Kan et al., 2008) also found that females were more vulnerable to ambient air pollution. Further research is needed to explore the source of sex differences in PM effects. In terms of age, we found generally more substantial impacts on children than older groups. Meanwhile these adverse effects diminished with age. Nevertheless, no significant differences between age groups were found, although children (Chen et al., 2019a; Chen et al., 2017; Zhang et al., 2019b) and the elderly (Hassanvand et al., 2017; Tian et al., 2019a) have been widely reported as vulnerable subpopulations for ambient particulate matters.

Besides, we observed significantly stronger PM-associated effects during the cold months, coincided with previous studies on ambulance emergency call-outs (Chen et al., 2020b), ED visits (Chen et al., 2019a) and hospital admissions for respiratory diseases (Zhang et al., 2019a). However, a study on ambulance emergency calls (Ai et al., 2019) found remarkably larger effects of PM<sub>2.5</sub> on all-cause and cardiovascular morbidity in the warm months, and slightly stronger effects on respiratory morbidity in the cold months. Furthermore, no substantially warm-cold differences were detected in another study on hourly pollutants level and the risk of myocardial infarction in England and Wales (Bhaskaran et al., 2011). These inconsistencies across studies might be attributable to discrepancies in regions, exposure patterns, pollution source and chemical compositions, as well as demographic characteristics (Shah et al., 2013; Zhang et al., 2019b). In addition, the exposure-response curves (Fig. 2) with a slower slope in the high concentrations

suggested that more severe PM emissions in the cold months could not be possibly responsible for the more harmful PM effects during the cold months. More sophisticated researches are needed to further clarify effects modification by warm-cold season.

Several limitations should be noted in this study. First, our exposure assessments of ambient PMs were averaged from the fixed stations, rather than individual measurements. This substitution may entail exposure misclassification (Zeger et al., 2000) and then underestimate the health effects of particulate matter (Li et al., 2019; Li et al., 2015). Second, our ED data were originated from one large-scale general hospital in Guangzhou city, so the results should be generalized to other regions with caution. Additionally, due to a lack of the corresponding classification of diseases, we were unable to investigate associations between size-specific PMs and cause-specific ED visits. Similarly, we cannot exclude ED visits caused by accident such as road traffic injuries and fires. Hence, we may underestimate the impacts of PMs on hourly ED visits to some extent (Cao et al., 2009).

## 5. Conclusions

In summary, we found very short-term impacts of ambient particulate matters on human health. Specifically, exposures to size-specific PMs may elevate risks of acute disease events in a few hours, and PM<sub>1</sub> entailed slightly stronger effects. In the cold months, these adverse effects were remarkably larger.

This study added to the evidence of PM<sub>1</sub> detrimental impact on human health at the hourly timescale, which may provide an informative reference in the establishment of hourly air quality standards and optimization of emergency medical resources. Additionally, this is a first study assessing detrimental effects of PM<sub>1</sub> of sub-daily exposures, thus the generalizability of our findings warrants further verification in the future research. To better alleviate disease burdens caused by ambient particulate matter, health policy makers and healthcare provider should take into consideration sub-micrometric and ultrafine particles, as well as finer temporal scales.

## CRedit authorship contribution statement

**Linjiong Liu:** Formal analysis, Writing - original draft, Writing - review & editing, Visualization. **Fujian Song:** Writing - review & editing.

**Jiaying Fang:** Resources. **Jing Wei:** Resources. **Hung Chak Ho:** Writing - review & editing. **Yimeng Song:** Resources. **Yuanyuan Zhang:** Data curation. **Lu Wang:** Data curation. **Zhiming Yang:** Writing - review & editing. **Chengyang Hu:** Writing - review & editing. **Yunquan Zhang:** Conceptualization, Methodology, Data curation, Writing - original draft, Formal analysis.

### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Acknowledgments

Yunquan Zhang was supported by the Key Research Center for Humanities and Social Sciences in Hubei Province (Hubei University of Medicine) (Grant No. 2020ZD001). Jiaying Fang was supported by the Health Science and Technology Project of Guangzhou (Grant No. 20191A011114). We acknowledge the China National Environmental Monitoring Center for providing air pollution data and United States' National Centers for Environmental Information for providing hourly meteorological data. We appreciated the anonymous reviewers very much, whose insightful comments and suggestions contributed a lot to improving the quality of our manuscript.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2020.142347>.

### References

- Ai, S., Wang, C., Qian, Z.M., Cui, Y., Liu, Y., Acharya, B.K., et al., 2019. Hourly associations between ambient air pollution and emergency ambulance calls in one central Chinese city: implications for hourly air quality standards. *Sci. Total Environ.* 696, 133956.
- Bhaskaran, K., Hajat, S., Armstrong, B., Haines, A., Herrett, E., Wilkinson, P., et al., 2011. The effects of hourly differences in air pollution on the risk of myocardial infarction: case crossover analysis of the MINAP database. *BMJ* 343, d5531.
- Bhaskaran, K., Armstrong, B., Hajat, S., Haines, A., Wilkinson, P., Smeeth, L., 2012. Heat and risk of myocardial infarction: hourly level case-crossover analysis of MINAP database. *BMJ* 345, e8050.
- Cao, J., Li, W., Tan, J., Song, W., Xu, X., Jiang, C., et al., 2009. Association of ambient air pollution with hospital outpatient and emergency room visits in Shanghai, China. *Sci. Total Environ.* 407, 5531–5536.
- Chen, G., Li, S., Zhang, Y., Zhang, W., Li, D., Wei, X., et al., 2017. Effects of ambient PM 1 air pollution on daily emergency hospital visits in China: an epidemiological study. *The Lancet Planetary Health* 1, e221–e229.
- Chen, G., Morawska, L., Zhang, W., Li, S., Cao, W., Ren, H., et al., 2018. Spatiotemporal variation of PM1 pollution in China. *Atmos. Environ.* 178, 198–205.
- Chen, D., Zhang, F., Yu, C., Jiao, A., Xiang, Q., Yu, Y., et al., 2019a. Hourly associations between exposure to ambient particulate matter and emergency department visits in an urban population of Shenzhen, China. *Atmos. Environ.* 209, 78–85.
- Chen, L., Zhang, Y., Zhang, W., Chen, G., Lu, P., Guo, Y., et al., 2019b. Short-term effect of PM1 on hospital admission for ischemic stroke: a multi-city case-crossover study in China. *Environ. Pollut.* 260, 113776.
- Chen, K., Schneider, A., Cyrus, J., Wolf, K., Meisinger, C., Heier, M., et al., 2020a. Hourly exposure to ultrafine particle metrics and the onset of myocardial infarction in Augsburg, Germany. *Environ. Health Perspect.* 128, 17003.
- Chen, T.T., Zhan, Z.Y., Yu, Y.M., Xu, L.J., Guan, Y., Ou, C.Q., 2020b. Effects of hourly levels of ambient air pollution on ambulance emergency call-outs in Shenzhen, China. *Environ. Sci. Pollut. Res. Int.* 27, 24880–24888.
- Di, Q., Dai, L., Wang, Y., Zanobetti, A., Choirat, C., Schwartz, J.D., et al., 2017. Association of short-term exposure to air pollution with mortality in older adults. *JAMA* 318, 2446–2456.
- Franck, U., Odeh, S., Wiedensohler, A., Wehner, B., Herbarth, O., 2011. The effect of particle size on cardiovascular disorders—the smaller the worse. *Sci. Total Environ.* 409, 4217–4221.
- Guo, Y., 2017. Hourly associations between heat and ambulance calls. *Environ. Pollut.* 220, 1424–1428.
- Guo, J.-P., Zhang, X.-Y., Che, H.-Z., Gong, S.-L., An, X., Cao, C.-X., et al., 2009. Correlation between PM concentrations and aerosol optical depth in eastern China. *Atmos. Environ.* 43, 5876–5886.
- Guo, J., Deng, M., Lee, S.S., Wang, F., Li, Z., Zhai, P., et al., 2016. Delaying precipitation and lightning by air pollution over the Pearl River Delta. Part I: observational analyses. *Journal of Geophysical Research: Atmospheres* 121, 6472–6488.
- Guo, J., Liu, H., Li, Z., Rosenfeld, D., Jiang, M., Xu, W., et al., 2018. Aerosol-induced changes in the vertical structure of precipitation: a perspective of TRMM precipitation radar. *Atmospheric Chemistry and Physics (Online)* 18.
- Hassanvand, M.S., Naddafi, K., Kashani, H., Faridi, S., Kunzli, N., Nabizadeh, R., et al., 2017. Short-term effects of particle size fractions on circulating biomarkers of inflammation in a panel of elderly subjects and healthy young adults. *Environ. Pollut.* 223, 695–704.
- Hu, K., Guo, Y., Hu, D., Du, R., Yang, X., Zhong, J., et al., 2018. Mortality burden attributable to PM1 in Zhejiang province, China. *Environ. Int.* 121, 515–522.
- Huang, K., Yang, X., Liang, F., Liu, F., Li, J., Xiao, Q., et al., 2019. Long-term exposure to fine particulate matter and hypertension incidence in China. *Hypertension* 73, 1195–1201.
- Hvidtfeldt, U.A., Geels, C., Sorensen, M., Ketznel, M., Khan, J., Tjonneland, A., et al., 2019. Long-term residential exposure to PM2.5 constituents and mortality in a Danish cohort. *Environ. Int.* 133, 105268.
- Institute, H.E., 2019. State of Global Air 2019. Health Effects Institute.
- Jiao, A., Xiang, Q., Ding, Z., Cao, J., Ho, H.C., Chen, D., et al., 2020. Short-term impacts of ambient fine particulate matter on emergency department visits: comparative analysis of three exposure metrics. *Chemosphere* 241, 125012.
- Kan, H., London, S.J., Chen, G., Zhang, Y., Song, G., Zhao, N., et al., 2008. Season, sex, age, and education as modifiers of the effects of outdoor air pollution on daily mortality in Shanghai, China: the public health and air pollution in Asia (PAPA) study. *Environ. Health Perspect.* 116, 1183–1188.
- Kim, J., Kim, H., Kweon, J., 2015. Hourly differences in air pollution on the risk of asthma exacerbation. *Environ. Pollut.* 203, 15–21.
- Le, C.T., Lindgren, B.L., 1988. Computational implementation of the conditional logistic regression model in the analysis of epidemiologic matched studies. *Comput. Biomed. Res.* 21, 48–52.
- Li, Z., Huang, G.H., Guo, L., Fan, Y.R., Chen, J.P., 2015. A fuzzy gradient chance-constrained evacuation model for managing risks of nuclear power plants under multiple uncertainties. *J. Environ. Inf.* 33 (2), 129–138.
- Li, Y.P., Huang, G.H., Cui, L., Liu, J., 2019. Mathematical modeling for identifying cost-effective policy of municipal solid waste management under uncertainty. *J. Environ. Inf.* 34 (1), 55–67.
- Lin, H., Tao, J., Du, Y., Liu, T., Qian, Z., Tian, L., et al., 2016. Particle size and chemical constituents of ambient particulate pollution associated with cardiovascular mortality in Guangzhou, China. *Environ. Pollut.* 208, 758–766.
- Lin, H., Liu, T., Xiao, J., Zeng, W., Guo, L., Li, X., et al., 2017a. Hourly peak PM2.5 concentration associated with increased cardiovascular mortality in Guangzhou, China. *J. Expo Sci Environ Epidemiol* 27, 333–338.
- Lin, H., Ratnapradipa, K., Wang, X., Zhang, Y., Xu, Y., Yao, Z., et al., 2017b. Hourly peak concentration measuring the PM 2.5-mortality association: results from six cities in the Pearl River Delta study. *Atmos. Environ.* 161, 27–33.
- Liu, H., Tian, Y., Xiang, X., Juan, J., Song, J., Cao, Y., et al., 2018. Ambient particulate matter concentrations and hospital admissions in 26 of China's largest cities: a case-crossover study. *Epidemiology* 29, 649–657.
- Liu, C., Chen, R., Sera, F., Vicedo-Cabrera, A.M., Guo, Y., Tong, S., et al., 2019. Ambient particulate air pollution and daily mortality in 652 cities. *N. Engl. J. Med.* 381, 705–715.
- Lumley, T., Levy, D., 2000. Bias in the case-crossover design: implications for studies of air pollution. *Environmetrics* 11, 689–704.
- Maclure, M., 1991. The case-crossover design: a method for studying transient effects on the risk of acute events. *Am. J. Epidemiol.* 133, 144–153.
- Phung, V.L.H., Ueda, K., Seposo, X., Takami, A., Sugata, S., Yoshino, A., et al., 2020. Hourly association between ambient PM2.5 and emergency ambulance dispatches in 11 cities in Japan. *Environ. Res.* 185, 109448.
- Qi, J., Ruan, Z., Qian, Z.M., Yin, P., Yang, Y., Acharya, B.K., et al., 2020. Potential gains in life expectancy by attaining daily ambient fine particulate matter pollution standards in mainland China: a modeling study based on nationwide data. *PLoS Med.* 17, e1003027.
- Shah, A.S.V., Langrish, J.P., Nair, H., McAllister, D.A., Hunter, A.L., Donaldson, K., et al., 2013. Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet* 382, 1039–1048.
- Tian, Y., Liu, H., Liang, T., Xiang, X., Li, M., Juan, J., et al., 2019a. Fine particulate air pollution and adult hospital admissions in 200 Chinese cities: a time-series analysis. *Int. J. Epidemiol.* 48, 1142–1151.
- Tian, Y., Liu, H., Wu, Y., Si, Y., Song, J., Cao, Y., et al., 2019b. Association between ambient fine particulate pollution and hospital admissions for cause specific cardiovascular disease: time series study in 184 major Chinese cities. *BMJ* 367, l6572.
- Wang, W., Mao, F., Zou, B., Guo, J., Wu, L., Pan, Z., et al., 2019. Two-stage model for estimating the spatiotemporal distribution of hourly PM1.0 concentrations over central and East China. *Sci. Total Environ.* 675, 658–666.
- Wei, J., Li, Z., Guo, J., Sun, L., Huang, W., Xue, W., et al., 2019a. Satellite-derived 1-km-resolution PM1 concentrations from 2014 to 2018 across China. *Environ Sci Technol* 53, 13265–13274.
- Wei, Y., Wang, Y., Di, Q., Choirat, C., Wang, Y., Koutrakis, P., et al., 2019b. Short term exposure to fine particulate matter and hospital admission risks and costs in the Medicare population: time stratified, case crossover study. *BMJ* 367, l6258.
- Wu, Q.Z., Li, S., Yang, B.Y., Bloom, M., Shi, Z., Knibbs, L., et al., 2020. Ambient airborne particulates of diameter  $\leq 1 \mu\text{m}$ , a leading contributor to the association between ambient airborne particulates of diameter  $\leq 2.5 \mu\text{m}$  and Children's blood pressure. *Hypertension* 75, 347–355.
- Yang, M., Chu, C., Bloom, M.S., Li, S., Chen, G., Heinrich, J., et al., 2018. Is smaller worse? New insights about associations of PM1 and respiratory health in children and adolescents. *Environ. Int.* 120, 516–524.

- Yang, B.Y., Guo, Y., Morawska, L., Bloom, M.S., Markevych, I., Heinrich, J., et al., 2019. Ambient PM<sub>1</sub> air pollution and cardiovascular disease prevalence: insights from the 33 communities Chinese health study. *Environ. Int.* 123, 310–317.
- Yin, P., Guo, J., Wang, L., Fan, W., Lu, F., Guo, M., et al., 2020. Higher risk of cardiovascular disease associated with smaller size-fractioned particulate matter. *Environmental Science & Technology Letters* 7, 95–101.
- Yorifuji, T., Suzuki, E., Kashima, S., 2014a. Cardiovascular emergency hospital visits and hourly changes in air pollution. *Stroke* 45, 1264–1268.
- Yorifuji, T., Suzuki, E., Kashima, S., 2014b. Hourly differences in air pollution and risk of respiratory disease in the elderly: a time-stratified case-crossover study. *Environ. Health* 13, 67.
- Zang, L., Mao, F., Guo, J., Gong, W., Wang, W., Pan, Z., 2018. Estimating hourly PM<sub>1</sub> concentrations from Himawari-8 aerosol optical depth in China. *Environ. Pollut.* 241, 654–663.
- Zang, L., Mao, F., Guo, J., Wang, W., Pan, Z., Shen, H., et al., 2019. Estimation of spatiotemporal PM<sub>1.0</sub> distributions in China by combining PM<sub>2.5</sub> observations with satellite aerosol optical depth. *Sci. Total Environ.* 658, 1256–1264.
- Zeger, S.L., Thomas, D., Dominici, F., Samet, J.M., Schwartz, J., Dockery, D., et al., 2000. Exposure measurement error in time-series studies of air pollution: concepts and consequences. *Environ. Health Perspect.* 108, 419–426.
- Zhang, Y., Ding, Z., Xiang, Q., Wang, W., Huang, L., Mao, F., 2019a. Short-term effects of ambient PM<sub>1</sub> and PM<sub>2.5</sub> air pollution on hospital admission for respiratory diseases: case-crossover evidence from Shenzhen, China. *Int. J. Hyg. Environ. Health* 113418.
- Zhang, Y., Fang, J., Mao, F., Ding, Z., Xiang, Q., Wang, W., 2019b. Age- and season-specific effects of ambient particles (PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub>) on daily emergency department visits among two Chinese metropolitan populations. *Chemosphere* 246, 125723.
- Zhang, Y., Xiang, Q., Yu, C., Yang, Z., 2020. Asthma mortality is triggered by short-term exposures to ambient air pollutants: evidence from a Chinese urban population. *Atmos. Environ.* 223.
- Zhou, M., Wang, H., Zeng, X., Yin, P., Zhu, J., Chen, W., et al., 2019. Mortality, morbidity, and risk factors in China and its provinces, 1990–2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 394, 1145–1158.